

## Isolation and study of a ubiquitously expressed tomato pectin methylesterase regulatory region

**Martín-Ernesto Tiznado-Hernández\***

Departamento de Tecnología de Alimentos de Origen Vegetal  
Centro de Investigación en Alimentación y Desarrollo, A.C.  
Carretera a la Victoria km. 0.6  
Apartado Postal 1735  
Hermosillo, Sonora, 83000, México  
Tel: 52 662 80 00 55  
Fax: 52 662 280 04 22  
E-mail: tiznado@cascabel.ciad.mx

**Joel Gaffe**

Genetique Moleculaire des Plantes  
Universite Joseph Fourier  
Cermo BP 53, 38041 Grenoble Cedex 9, France  
Fax: 33 4 76 51 43 36  
Tel: 33 4 76 51 44 41  
E-mail: joel.gaffe@ujf-grenoble.fr

**Avtar K. Handa**

Department of Horticulture and Landscape Architecture  
PURDUE University  
1165 Horticulture Building  
West Lafayette, IN, 47907-1165, USA  
Tel: 765 494-1339  
Fax: 765 494-0391  
E-mail: handa@hort.purdue.edu

**Keywords:** pectin Methylesterase, promoter analysis, tobacco transgenic plants, tomato.

**Pectin methylesterase (PME) is an enzyme located in the plant cell wall of higher plants whose physiological role is largely unknown. We had isolated a PME gene from a tomato genomic library, including 2.59 kb of 5' flanking region and the coding region. Both coding and promoter region were sequenced and computer analyzed. Tobacco transgenic plants were created harboring constructs in which 2.596 Kb, 1.306 Kb and 0.267 Kb sizes of the promoter were driving the expression of  $\beta$ -Glucuronidase gene (GUS). GUS activity was studied by histochemical and fluorometric assays. Two introns of 106 and 1039 bp were found in the coding region and phylogenetic analysis placed this PME gene closer to genes from *Citrus sinensis* and *Arabidopsis thaliana* than tomato fruit-specific PME genes. In the promoter, it was found direct repeats, perfect inverted repeats and light responsive elements. GUS histochemical analysis showed activity in all plant tissues with the exception of pollen. The reduction in the promoter size induced a reduction in GUS activity in root, stem and leaf. Furthermore, root and leaf showed the highest and lowest activity, respectively. We had isolated a tomato PME gene with novel characteristics as compared with other known PME genes from tomato.**

Pectin methylesterase (PME) is an enzyme that have been found in every plant tissue analyzed (Lineweaver and Jansen, 1951; Rexova-Benkova and Markovic, 1976), several fungi (Christgau, et al. 1996; Mendgen, et al. 1996), bacteria (Plastow, 1988; Barras et al. 1994) and even insects (Ma et al. 1990; Shen et al. 1999). In higher plants, it is known to be a cell wall associated protein and several of the PME cDNA available in the literature, are known to have toward the N-terminal sequence, a characteristic signal peptide which is thought to help in targeting the protein to the plant cell wall (Gaffe et al. 1997 and references therein). PME catalyzes the deesterification of galactosyluronate methylesters of pectins, releasing protons and methanol into the media (Frenkel et al. 1998). Despite the biochemical mode of action of PME is well known, it have been difficult to demonstrate any role for PME in the physiology of plants. However, several hypothesis had been proposed: pollen germination and/or tube growth (Mu et al. 1994), abscission (Sexton and Roberts, 1982), regulation of cell enlargement through changes in the plant cell wall Donnan potential (Ricard and Noat, 1986), fruit softening during postharvest fruit ripening (Zeng et al. 1996) and plant defense (Chamberland et al. 1991; Wietholter et al. 2003). Furthermore, strong experimental evidences had

\*Corresponding author

been provided to suggest that a PME role in the release of cells from the root cap (Stephenson and Hawes, 1994; Wen et al. 1999), plant pathogenesis (Collmer and Keen, 1986; Mendgen, et al. 1996; Nun et al. 1996; Valette-Collet et al. 2003), plant systemic infection by tobacco mosaic virus (Chen and Citovsky, 2003) and maintenance of the tomato fruit tissue integrity during postharvest shelf life (Tieman and Handa, 1994). However, the actual physiological role of PME is still matter of controversy.

Four papers to our knowledge had been published in which the cloning of regulatory sequences of PME from higher plants was described. However, in all of them DNA comparison by computer was the only tool used to prove that the gene located downstream was indeed encoding a pectin methylesterase. Albani et al (1991) reported the finding of a genomic clone from *Brassica napus* which contains the PME gene and its 5' upstream regulatory region. Studies were conducted using a piece of the gene located downstream as a probe. This gene was found to be expressed mainly during pollen development. Two putative PME promoter regions were cloned from *Brassica campestris* (Kim et al. 1997). Study of their sequence found them to have high homology with the previously reported promoter PME from *Brassicca napus* (Albani et al. 1991). Further, a sequence motif similar to the one known to exist in two tomato pollen-specific promoter was located. Tobacco transgenic plants with constructs containing two different promoter sizes from one of those two promoter available were made. Expression of the GUS gene was only detected during developing and mature pollen grains germinated in vitro. Recently, the cloning of two 5' upstream region of PME from *Citrus sinensis* was published. Northern blot analysis showed that both DNA regulatory region are active in most of vegetative tissues (Nairn et al. 1998).

In our laboratory, we have cloned a 13.7 kb. genomic DNA from tomato containing a 2.59 kb. of DNA 5' flanking region, along with all the PME genomic clone. Identification of the protein encoded by the gene downstream was made by creating tobacco transgenic plants over expressing the PME cDNA (Gaffe et al. 1997) and comparing the sequences of the genomic and cDNA regions. In this work, we describe the study of a 5' flanking region of a PME gene called pmeu1 (which stands for PME ubiquitous one) using computer tools and tobacco transgenic plants.

## MATERIALS AND METHODS

### Cloning of the genomic fragment

The plaque lift technique was used to screen 810,000 clones of a tomato cv 'Cherry' genomic library made in EMBL3A  $\lambda$  phage with a radiolabeled small piece of PME; cloned by RT-PCR from tomato roots poly-A<sup>+</sup> mRNA (Gaffe et al. 1995). We found several hybridizing, putatively positives, plaques. From every plate, we made two lifts and only that

plaques producing signal in both lifts were chose to continue. After four rounds of purification and screening, one plaque turns out to be positive. Elimination of the bacteria present in the agar was made by using chloroform-containing SM buffer. For phage amplification, *E. coli* strain LE 392 was infected with the phage after cultured in LB media. DNA isolation from the phage was made through a phenol chloroform protocol (Ausubel et al. 1988). Digested DNA with several restriction enzyme was separated by electrophoresis and blotted into nylon membranes. These DNA blots were probed with the PME cDNA complete sequence available to locate the phage DNA region encoding the genomic PME gene and the region 5' upstream. Sal I and EcoR I digested DNA fragments were subcloned into pBSKS (+/-) vector (STRATAGENE CLONING SYSTEMS. La Jolla, CA).

The EMBL3A  $\lambda$  phage library screened was created using the Sal I restriction site. Because of this, digested DNA fragments using Sal I were used to calculate the size of the tomato DNA inserted into the phage isolated, found to be 13.7 kb. All the procedure above mentioned was performed essentially as described (Sambrook et al. 1989) unless otherwise indicated.

### DNA sequencing of the promoter and the genomic coding region

Nested unidirectional deletions of the 5' upstream DNA Sal I fragment were made by following the recommendations of the company (Erase-a-Base<sup>®</sup> System, Promega Corporation, Madison, WI). Deleted clones with about 250 bp of size difference were used for DNA sequencing using the T<sub>3</sub> universal primer by the Sanger dideoxy chain termination technique following the recommendations (Sequenase Kit, United States Biochemical Corporation, Cleveland, Ohio). Second strand sequencing was determined by the DNA sequencing facility of IOWA State University by using primers designed at proper positions in the sequence (Iowa State University, Ames, Iowa).

### Creation of the constructs

Three chimeric constructs driving the  $\beta$ -glucuronidase gene (*uidA*) under different sizes (2.596 Kb, 1.306 Kb and 0.267 Kb) of the promoter region were created by transcription fusion through the insertion of two stop codons in between the ATG of the pmeu1 gene and the ATG of the *uidA* gene. Every chimeric construct was ligated into the promoterless binary vector pBI101.3 (Bevan, 1984). This plasmid includes the neomycin phosphotransferase gene (NPTII) which confers resistance to kanamycin to be used as selectable marker. Furthermore, in this plasmid the DNA introduced is located between the right and left borders of the T-DNA, which allows the transference into the plant genome by *Agrobacterium* infection (Hooykaas, 1989; Zupan and Zambryski, 1995; Nester et al. 1996). Proper insertion of the different promoter sequences into the plasmid was confirmed by DNA digestion using suitable

restriction enzymes and PCR using primers designed against sequences in the PME promoter and *uidA* gene.

Every chimeric construct created included 150 bp in between the ATG of the *pmeu1* gene and the ATG of the *uidA* gene, containing sequences from the *pmeu1* gene and pBSKS(+/-) phagemid and pBI101.3 binary vector. Sequencing between the two ATG's was used to verify the presence of two stop codons and to corroborate the transcription fusion of the two ATG.

### Tobacco transgenic plants

Mobilization of the pBI101.3 plasmid into *Agrobacterium* LBA4404 was performed by triparental mating using the broad-host helper plasmid pRK2013 (Ditta, 1981). *Agrobacterium* transconjugants were screened on plates containing a mixture of kanamycin and rifampicin antibiotics in YEP media (Sambrook et al. 1989). Verification of the mobilization of the constructs was made by purification DNA from *Agrobacterium* following the recommendations (Wizard Minipreps, Promega Corporation, Madison, WI) and digestion with proper restriction enzymes. Tobacco (*Nicotiana tabaccum* W38) young leaves were infected with *Agrobacterium* by using the leaf disk technique (Mathis and Hichee, 1994) and selection for transformants was done by using kanamycin in the media.

### GUS activity measurement

After induction of roots, about 50 primary independent transformant plants growing *in vitro* harboring every of the three constructs were selected at random to measure GUS activity in leaf. This was done using the fluorometric technique (Jefferson et al. 1987) with a Perkin Elmer LS5 fluorometer. Quantification of reaction product was done by using a 4-methylumbelliferone standard curve. Also, six independent transgenic plants were used to measure GUS in root, stem and leaf tissues. Every GUS measurement was done at least three times. For enzymatic specific activity, protein determination was made using Bradford (1976) with bovine serum albumin as standard.

In order to examine the GUS presence in different tissues, at least 20 primary transgenic plants harboring the different constructs, were vacuum infiltrated with a 1.9  $\mu$ M solution of 5-Bromo-4-Chloro-3-Indolyl-Glucuronide (Jefferson et al. 1987) as described (Mandel et al. 1995).

### Pollen germination *in vitro*

Tobacco flowers in anthesis were collected from plants growing in the greenhouse and transported immediately to the laboratory. Anthers were cut and only that pollen released by a gently shaking was used for germination studies. Pollen was germinated using the Brewbaker and Kwack solution as described (Brewbaker and Kwack, 1963). Histochemical GUS staining was performed after four hours of pollen germination. Germination solution was

changed by the GUS staining solution and left at 37°C for at least 18 hrs before examination for GUS staining.

### Computer analysis

DNA sequence from the different *pmeu1* 5' flanking region deletions were joined together using DNAsis (Hitachi Software Engineering Co., LTD., 1991). Comparison between the *pmeu1* cDNA and *pmeu1* genomic clone was performed using Harr plot analysis with DNAsis software. Presence of known cis-acting elements was determined using the programs MathInspector ver. 2.2 (Quandt et al. 1995), TFSEARCH ver. 1.3 (Parallel Application, Tsukuba Laboratory, RWCP, Japan), Signal Scan ver. 4.05 (Prestridge, 1991) and Pattern Search (Wingender et al. 1996; Wingender et al. 1997). Percent of identity among the different PME promoters and PME transcribed regions were determined using Align (Myers and Miller, 1988). Alignment of deduced amino acid sequences was performed using GCG's Pileup Program (Genetics Computer Group, Madison, WI). Multiple sequence alignment was performed using CLUSTAL W (Thompson et al. 1994). DNA direct repeats for the tomato PMEUI promoter were determined using Proscan ver 1.7 and repeats from GCG software ver. 9.0 (Genetics Computer Group, 1995). Perfect inverted repeats (mirror repeats) were located using Palindrome from GCG software ver 9.0 (Genetics Computer Group, 1995). Putative TATA box was located by Signal Scan ver. 4.05. Phylogenetic analysis were done using the phylogeny inference package (Felsenstein, 1989; Felsenstein, 1993).

### Statistical analysis

Comparison of leaf GUS activities for the three constructs and for the different tissues was made by variance analysis using a completely randomized design for unbalanced number of repetitions. Tukey test was used when needed to find differences among means. Because it is known that the GUS enzymatic activity in populations of first-generation transgenic plants does not follow a normal distribution (Nap et al. 1993), we performed a Box-Cox transformation before variance analysis. From here, we learned that a square root was a suitable transformation to bring the GUS activity parameter into normality. Statistics reported in this paper represents the back transformation of the square root transformed data. All statistical analysis were performed using the SAS software (SAS Institute Inc. Cary, N.C.).

## RESULTS

### Isolation and characterization of PMEUI gene

The cloning and characterization of the entire PMEUI tomato cDNA has been previously reported (Gaffe et al. 1996; Gaffe et al. 1997). The next step lead us to the isolation and characterization of the genomic fragment

containing the PMEU1 gene. An EMBL3A phage of a tomato genomic library (VNTF cherry) was screened using 300 bp cDNA fragment corresponding to the conserved PME domain in PMEU1 (Gaffe et al. 1996; Gaffe et al. 1997). Four rounds of phage amplification allowed us to purify a single positive clone.

Subcloning, analysis by restriction mapping and DNA blot of the tomato genomic DNA fragment contained in the EMBL3A phage indicated that the size of the inserted tomato genomic DNA is 13.7 kb and the PMEU1 gene was found to be located toward the 5' region, spanning 5.28 kb.

In [Figure 1](#), is presented the organization of the EMBL3A clone containing the PMEU1 gene. This region includes 2.59 kb of DNA regulatory region and 2.89 kb of DNA transcribed region, shown as white and black areas. In the figure it is also shown the location of the right and left lambda phage arms and the main restriction sites.

### DNA sequence of the transcribed region of PMEU1 gene

In [Figure 2](#), it is shown the sequence of the PMEU1 genomic clone (GenBank Accession Number: AY046596). In italics, it is presented the 5' untranslated region (Gaffe et al. 1997) and the partial 3' untranslated region. In bold, it is shown the sequence of the two introns present. Underlined, it is presented the translation start site and stop codon (TAA). Double underlined it is shown the putative polyadenylation signal and polyadenylation site (GT).

The polyadenylation signal was found to follow the plant consensus sequence AAUAAA (Li and Hunt, 1995). The two introns present are of 106 and 1039 bp in length. Both of them showed a significantly higher composition of U's with respect to the flanking exon sequences. This is a characteristic known to be present in many plant genes (Ko et al. 1998).

### Intron-exon organization of PMEU1 and other PME genomic clones

The intron-exon structure of the PME genomic sequences available has been analyzed. The splice junction of all the clones conform to the GT/AG boundary rule for the 5' donor and 3' acceptor site (Liu and Filipowicz, 1996). The intron size range from 72 to 1577 bp and the exon from 117 to 1353 bp. The average value for intron and exon size is 109 and 519, respectively.

Seventeen clones have only one or two introns. Three putative PME genomic sequences from *Arabidopsis* contains four introns and show a level of similarity with PMEU1 of around 50%. Further, AtPME7 with five introns is more closely related to PMEU1 (64.9% of similarity). These observations suggest that there is not a simple relationship between the phylogenetic distance and intron number in the different PME genomic clones.

The position of one intron, relative to the deduced amino acid sequence, is conserved in 19 out of the 22 plant PME genomic sequences. This intron is located 17 amino acid residues upstream of the PME signature sequence GPXKHQAVLR; observed in the rice genomic clone as well ([Figure 3](#)). This observation suggests that monocots and dicots share a common ancestor. The other three clones (AtPME8, AtPME9 and AtPME10) are clustered together in one group by the phylogenetic analysis ([Figure 4](#)) which agrees with the lack of the intron located at the same distance from the signature sequence and the common characteristic of the presence of four introns.

### Phylogeny analysis among PMEU1 and other plant PME genes

Deduced amino acid sequences of 22 plant PME genes as well as PMEB from *Erwinia chrysanthemi* were included in our study. The plant PME genes were chosen based in published data providing experimental evidences or presence of the full genomic sequence from the *Arabidopsis thaliana* genome project from which some of the PME genes were included. One of the pectin methylesterase genes from *Oriza sativa* was included to be able to compare with a PME from monocots. Furthermore, the gene from *E. chrysanthemi* was chosen in order to compare PME from plants with a distantly related PME and also to have a control in the phylogenetic analysis. The PMEU1 gene includes 2900 bp and a theoretically deduced open reading frame of 583 amino acids ([Figure 3](#)). Several sequences shorter than 400 bp like PECS-1-2 from *Citrus sinensis*, are known to be partial. However, PPE1 sequence from *Petunia inflata* is shorter than 400 bp and still encodes a full polypeptide.

Sequence alignment of these different encoded polypeptides indicate that the N-terminal half of these clones is loosely conserved compared with the C-terminal half, involved perhaps in the PME catalytic activity ([Figure 3](#)). Because of this, a final alignment, edited to represent only the phylogenetically relevant fraction of the sequences was used to derive a phylogenetic tree ([Figure 4](#)).

Based on this phylogenetic analysis, we organized up to 18 genomic clones in five groups. Five PME genomic clones from various origins can not be associated with any of these groups. The lack of association of PME from *Erwinia chrysanthemi* with other plant PME's was something expected, however, it is interesting that the clone PECS-2.1 from *Citrus sinensis* is distantly related with the two clones PECS-1.1 and PECS-1.2 from the same source that clustered together with the PMEU1 clone.

This phylogenetic analysis indicates that PMEU1 belong to a group containing two *Citrus sinensis* PME genes, PECS-1.1 and PECS-1.2 and two *Arabidopsis thaliana* genes, AtPME2 and AtPME3; however, it is distant from the three tomato PME genes expressed only in tomato fruit tissues: LePME1, LePME2 and LePME3 (Harriman et al. 1992),

suggesting the PMEU1 is a gene evolved to have a different and novel function. However, due to the limited amount of information concerning the expression of these genes, we can not establish a clear relationship between these groups of PME genes and their possible function.

### Structure of PMEU1 promoter

In [Figure 5](#) it is shown the 2.59 kb. PMEU1 promoter sequence (GenBank Accession Number AY050764). Computer study of this sequence showed several features commonly present in DNA regulatory sequences. The largest direct repeats within the promoter sequence, are shown underlined and numbered. Mirror repeats are shown with arrows in opposite directions. Putative cis-acting elements are shown boxed and roman numbered. The putative TATA box is shown double underlined. In bold, it is shown the transcription start site.

Study of the 5' region of this sequence did not indicate the presence of elements commonly present in the 3' region of genes, suggesting that the PMEU1 promoter region could be larger than 2.59 kb.

The number of direct repeats located by computer in the PMEU1 promoter varied with the size of the fragment, in such a way that it was found only one for repeats consisting of 17 and 26 bp, four for repeats with 12 bp, three for repeats with 11 bp and greater than 1000 for repeats with 5 bp (data not shown). However, the significance of this repeats within the PMEU1 promoter remains to be elucidated.

We also locate in the promoter sequence several perfect inverted repeats or mirror repeats, depicted in [Figure 5](#) as arrows pointing in opposite directions. It is interesting that the longest inverted repeats is contained within the longest direct repeats. As in the case of the direct repeats, the function of these inverted repeats, if any, is unknown.

Short sequences with resemblance to known cis-acting elements present in other ADN regulatory regions were located in the PMEU1 promoter sequence. In [Figure 6](#) are included only the ones with the highest degree of similarity. Two copies of the sequence GAAAGA shown to confer responsiveness to red light in the phytochrome A3 promoter (Bruce et al. 1991) are present in PMEU1 promoter (box I). Also, one copy similar to the sequence GTGAGGTAATAT, known to be regulated by light (Fluhr and Dankekar, 1986; Green et al. 1987) was found (box II). Furthermore, we found regions similar to a G-box (box III), shown to be light inducible (Schindler et al. 1992). Also, it was located a sequence similar to an abscisic acid responsive element (box IV) (Guiltinan et al. 1990). As can be seen from above, three of the four putative cis-acting elements located are known to be regulated by light. Experiments to show whether PMEU1 promoter is regulated by light deserves further attention. However, still the function of this cis-acting elements within the PMEU1

promoter is largely theoretical and experimental evidences to confirm any function of these sequences remains to be provided.

We were able to locate a putative TATA box 44 bp upstream of the transcription start site ([Figure 5](#)). However, as mentioned for the other elements above described, the confirmation of this region as actual TATA box still need to be experimentally probed. We did not find the presence of a CAAT box, although it had been shown to be present in several promoter of plant genes (Joshi, 1987).

Paired comparisons among the DNA sequence of the PMEU1 promoter with sequences of PME promoters from *Brassica campestris* (GBAN215-6 and GBAN215-12), *Brassica napus* (Bp 19), *Citrus sinensis* (CsPME1 and CsPME3) and *Arabidopsis thaliana* (AtPME1) did not showed any special pattern or similarity with any of the promoters included in the analysis. Indeed, all the pair comparisons showed around 50% of identity. Further, analysis by multiple sequences alignment among all PME promoters failed to locate an homologous region in common to all of them (data not shown).

### Transgenic tobacco plants

With the goal to test whether the 2.59 kb. DNA region located in the 5' flanking region of the PMEU1 genomic coding region represent an active promoter, we created several tobacco transgenic plants expressing chimeric constructs in which 2.59, 1.3 and 0.267 kb of promoter sequence is driving the expression of the reporter gene *uidA* encoding the  $\beta$ -glucuronidase enzyme.

In [Figure 6](#) it is shown the three constructs made along with the average of leaf GUS activity for about 50 independently tobacco transformed plants growing *in vitro* and expressing the corresponding construct. From the graph, it is clear the trend: the bigger the piece of the promoter, the higher the activity of the *uidA* gene. Statistical analysis of root squared-transformed data found differences among all of them ( $p < 0.05$ ).

Histochemical staining of many independent primary tobacco transgenic seedlings showed activity in leaf, stem and roots of the plants. We also found activity in petals and sepals. However, no activity was detected in pollen grain or *in vitro* germinated pollen (data not shown).

In [Figure 7](#), it is shown the average values of GUS activity for root, stem and leaf of six independent tobacco plants harboring every of the three constructs. The effect of reducing the size in the PMEU1 promoter for the different tissues analyzed followed the pattern already observed in leaf. The decrease in the size of the PMEU1 promoter region reduce its transcriptional activity in all differentiated tissue analyzed.

Statistical analysis found significant differences ( $p < 0.05$ )

among the root tissues from plants harboring the different sizes of the promoter. For stem tissues, significant differences were found only between plants with 0.267 kb and 2.59 kb of promoter size. This result is most likely due to the few independent transformants used in the analysis. However, the trend is clear and similar in all plant differentiated tissues analyzed.

## DISCUSSION

We have cloned and analyzed a genomic DNA region containing an almost complete and novel PME gene. Several tools were used to probe that this region encodes the genomic sequence of a PME gene. Comparison of the sequences of PMEU1 genomic coding region with the PMEU1 cDNA already cloned showed that both are identical with the exception of the intron sequences located in the genomic clone. Further, analysis of the cDNA sequence using BLAST resulted in high similarity with several DNA regions encoding PME genes. Also, transgenic plant overexpressing the PMEU1 cDNA under the control of the cauliflower mosaic virus showed higher levels of PME activity as compared with control plants. It was also shown that this high level of PME activity correlated with the presence of a band hybridizing with a PMEU1 specific probe (Gaffe et al. 1997).

The PMEU1 gene is presented in the tomato genome as a single copy (Gaffe et al. 1997), in contrast with other PME genes published which had been shown to form clusters (Richard et al. 1996; Turner et al. 1996).

We perform several experiments to find another copy of the gene, like increasing the number of plaques screened and using probes from the 5' end of the gene with unsuccessful results. Also, DNA blot analysis of the 8.4 kb of the 3' end of the DNA inserted in the phage did not show any hybridization with PMEU1 probe even under low stringency conditions (data not shown). Further, DNA blot analysis of the tomato genome using EcoR I as restriction enzyme showed one band hybridizing to a 6.0 kb band, which correspond precisely with the fragment released from the DNA phage and shown to hybridize with the PMEU1 specific probe (data not shown). Taken together, these evidences support that the PMEU1 gene is presented as a single copy in the tomato genome and that it is part of the DNA contained by the isolated phage from the genomic library.

Comparison of the PMEU1 genomic coding region with the PMEU1 cDNA sequence showed the presence of two introns with 106 and 1039 bp in size (Figure 2). We compared the structure of genomic regions encoding PME genes in regard to the number and size of introns. The analysis did not show any clear pattern of structure since there is a high variability in both the size and the number of introns present. However, when we compared the amino acid sequence of 23 PME genes from higher plants and a PME gene from *E. chrysanthemi* (Figure 3), the analysis highlighted a large region in common for most of the plant

PME genes: GPXKHQAVALLR. Also, we noted that it is located most of the time at the same place with respect to the presence of the first intron. Experiments of site directed mutagenesis with a PME gene from *Aspergillus niger* strain 5344 had shown that there is an histidine residue essential for PME activity within the amino acid sequence HQAVA (Duwe and Khanh, 1996). From Figure 3, we can see that most of the PME enzymes from higher plants has the sequence HQAVA as well. This seems to suggest that this histidine residue can be playing an important role in the catalytic activity of the enzyme. Multiple sequence alignment failed to locate the sequence of HQAVA of *Erwinia chrysanthemi* PME A or PMEB at the same location as plant PME's. However, pair comparison between PMEU1 and PME A or PMEB from *Erwinia chrysanthemi* correctly aligned the sequence HQAVA at the same position.

Studies of the three-dimensional structure of *Erwinia chrysanthemi* pectin methylesterase (PME-A) support the presence of two aspartate and one arginine residues in the active site of the enzyme (Jenkins et al. 2001) and not an histidine. However, some of the PME isoenzymes show an aspartate residue instead of histidine in the same site (Figure 3).

We believe that the study of the possible involvement of either an histidine or an aspartate residues in the catalytic activity of PME from higher plants deserves further attention.

Computer analysis of the PMEU1 genomic region showed that this sequence follows several features commonly present in other genes from higher eukaryotic organisms, as mentioned above. The phylogenetic analysis (Figure 4) had shown that this PME gene is not related with other PME genes isolated from the tomato genome (Harriman et al. 1991). Rather, from Figure 4, we can see that PMEU1 is more related to two genes from *Arabidopsis thaliana* (AtPME2 and AtPME3) and two genes from *Citrus sinensis* (PECS-1.1 and PECS-1.2). Efforts to find a correlation between relatedness of the PME genes and pattern of expression were not successful. However, the finding just mentioned further support that the cloned PME gene described in this work belong to a entirely novel type of PME gene from tomato.

Experiments carried out in our lab with tobacco transgenic plants overexpressing the PMEU1 gene and tomato plant with lower levels of this gene did not produce a change in the plant phenotype that could be give us an insight as to what is the physiological role of the PMEU1 gene. Therefore, we decided to computer analyzed the PMEU1 promoter sequence to look for DNA boxes or elements with known function, in search for insights as to what can be the physiological role of this PMEU1 gene.

In Figure 5, it is presented the sequence of the DNA regulatory region of the PMEU1 gene. We are not sure of

having the complete genomic sequence of the PMEUI gene for two reasons: the DNA segment of the PMEUI gene was located toward the 5' end of the tomato genomic DNA carried by the isolated phage (Figure 1). Further, computer analysis of the PMEUI promoter 5' end region failed to find elements known to exist toward the 3' end of the gene coding regions. However, considering the size of the largest sequence of a PME regulatory region published to date, 2.3 kb (Albani et al. 1991), it is quite possible that we almost had the entire PMEUI regulatory region. Our efforts to isolate from the tomato genomic library the remaining segment of the PMEUI regulatory region were largely unsuccessful.

The computer analysis of the PMEUI regulatory region showed the presence of both direct repeats and perfect inverted repeats. In Figure 5, only the largest ones are shown. It is interesting that repeats 1 and 2, which are only separated by one base pairs appears to come from only one repeat in which a mutation took place, splitting this long repeats into two shorter ones. Also, some of the largest perfect inverted repeats are present inside of the largest direct repeats. It can be interested to test whether this repeats belong to the PMEUI promoter or they are part of the intergenic region of the plant genome which is known to contain repeat sequences. However, the possible role if any of these repeats remains to be elucidated.

We also located two sequences identical to cis-acting elements found in the phytochrome A3 promoter (Bruce et al. 1991). Also, it showed two more sequences similar to known cis-acting elements regulated by light. From here, the possible regulation of this PMEUI gene by light deserves further attention. We also located a sequence similar to a known abscisic acid responsive element, close to the transcription start site (Figure 5). The phytohormone ABA had been related to the abscission phenomena in plants (Label et al. 1994; Aneju et al. 1999) and to the plant responses to abiotic stress in plant (Zhu, 2001). One of the genes encoding a pectin methylesterase isolated from *Citrus sinensis* was shown to be up-regulated in abscission zones of leaves (Nairn et al. 1998). Currently, experiments in our laboratory are being carried out to test the possible role of the gene PMEUI in the plant responses to light, abscission and abiotic stress, however, a possible function for the PMEUI gene in these phenomena is still matter of controversy.

With the goal to demonstrate that the 5' flanking region of the PMEUI genomic clone correspond with an active regulatory region, and to find the smallest size of the region able to direct transcription, we created transgenic tobacco plants expressing different constructs in which the *uidA* gene, encoding the enzyme  $\beta$ -glucuronidase, is being regulated by different regions of the PMEUI promoter.

In Figure 6, it is shown the results of analyzing the  $\beta$ -glucuronidase activity of around 50 independent transformed tobacco plants. From the figure, it is clear that

by reducing the size of the promoter, its transcriptional activity is also reduced. As can be seen, even 267 bp of the PMEUI regulatory region is transcriptionally active. This means that we did not reach the lower limit where the promoter loose completely its transcriptional activity, although a large reduction was accomplished. In contrast, it was reported that a truncated piece of 440 bp of a flax PME promoter (Lupme3) lost completely the ability to drive transcription of a reporter gene (Roger et al. 2001). The results of GUS activity in leaf tissue are supported by the histochemical staining analysis in which the transgenic plants showed weaker activity in the parenchyma tissue surrounding the leaf vascular tissue with decrease in the promoter size (data not shown).

The change in transcriptional activity among the different sizes of promoter is of 6 fold when comparing the 0.267 kb. with the 1.306 kb. and 4 fold when comparing the 1.306 kb. with the 2.59 kb. There is a difference of 1.03 kb between 0.267 kb and 1.306 kb and 1.29 kb between 1.306 kb and 2.59 kb. The differences in sizes are similar and still the variation in activity is higher between the 0.267 Kb. and 1.306 kb which means that perhaps there are stronger enhancer element(s) in the promoter region closest to the ATG. Overall, we obtained up to 95% in reduction of PMEUI promoter transcriptional activity with the construct including 0.267 kb of PMEUI promoter. Reduction of the promoter size which brings an associated reduction in promoter activity as measured with a reporter gene had been found in deletion studies of other promoters (Darasiela et al. 1996; Royo et al. 1996), however, sometimes smaller pieces are able to drive higher levels of reporter gene activity in general (Canevascini et al. 1996) or at some specific tissues (Royo et al. 1996).

The standard deviation of the parameter is indicating a very high variability which is most likely due to the presence of multiple copies in the genome of the different transformants (not determined), dissimilarities in the physiological status among the leaf tissues used and to the position effect (Wilson et al. 1990). This result is alike with studies reported earlier, in which a high variability among independent transformants was also found in liquid cell cultures expressing the GUS gene under the manopine synthase (Peach and Velten, 1991). Also, tobacco cells stably transformed with a chimeric construct in which the CaMV35S was driving the expression of GUS, showed a standard deviation three times higher than average for the GUS specific activity parameter (Allen et al. 1993).

The average of GUS activity in tobacco leaf for the construct harboring the 2.59 kb of promoter size was 324.334 pMoles of MU/min/mg protein (Figure 7). This activity is similar to the one reported earlier for tobacco (*Nicotiana tabacum* var Samsun) leaf of about the same size used in this work, harboring GUS (*uidA* gene) under the control of the cauliflower mosaic virus 35S promoter: 321 pMoles/min/mg protein (Jefferson et al. 1987). This result suggest that PMEUI promoter is as strong as the

CaMV35S which in turn indicates its usefulness in overexpressing proteins in plants.

We also studied the expression of the three constructs in the three main plant tissues: root, stem and leaf of six independent transformants (Figure 7). It is clear from the graph that the three constructs showed the same pattern already observed for leaf tissue. However, we recorded 1.7 and 8 fold PMEUI transcriptional activity for stem and root, respectively. This suggest a difference in the strength of the enhancer elements present in the PMEUI promoter depending upon the type of plant tissue. These results also suggest that the enhancer element(s) are active en several differentiated tissues and are not specific for leaf tissue. These results are supported by the GUS histochemical staining in which the transgenic plants harboring the construct including the smallest promoter region showed weaker activity in the parenchyma tissue surrounding the vascular tissue as compared with tissues of transgenic plants expressing the construct with the highest promoter region (data not shown).

These findings are in contrast with deletion studies of other promoter in which it was found that for specific tissues, smaller pieces of the regulatory regions are able to direct higher values of reporter gene enzymatic activity (Royo et al. 1996).

Deletion studies of the PMEUI promoter could be of significant insight to locate this putative enhancer elements. However, stronger experimental evidences are needed to probe their presence in the PMEUI promoter region.

In summary, we had isolated an entirely new gene encoding a pectin methylesterase isozyme from the tomato genome which is represented by a single copy. It shows an ubiquitous pattern of expression, in contrast with the tissue specific gene isolated earlier from tomato. Analysis of its promoter region suggest several potential function for this gene and we believe that further analysis of this gene will bring new insights to understand better the physiological role of the pectin methylesterase enzyme.

## REFERENCES

ALBANI, D.; ALTOSAAR, I.; ARNISON, D.G. and FABIJANSKI, S.F. A gene showing sequence similarity to pectin esterase is specifically expressed in developing pollen of *Brassica napus*. Sequences in its 5' Flanking Region are Conserved in Other Pollen-Specific Promoters. *Plant Molecular Biology*, April 1991, vol. 16, no. 4, p. 501-513.

ALLEN, G.C.; HALL, G.E.; CHILDS, L.C.; WEISSINGER, A.K.; SPIKER, S. and THOMPSON, W.F. Scaffold attachment regions increase reporter gene expression in stably transformed plant cells. *Plant Cell*, June 1993, vol. 5, no. 6, p. 603-613.

ANEJU, M.; GIANFAGNA, T. and NG, E. The roles of abscisic acid and ethylene in the abscission and senescence of cocoa flowers. *Plant Growth Regulation*, March 1999, vol. 27, no. 3, p. 149-155.

AUSUBEL, F.M.; BRENT, R.; KINGSTON, R.E.; MOORE, D.D.; SEIDMAN, J.G.; SMITH, J.A. and STRUHL, K. *Current Protocols in Molecular Biology: Screening of Recombinant DNA Libraries*. John Wiley & Sons, Inc. 1988. p. 1600. ISBN 047150338X.

BARRAS, F.; van GIJSEGEM, F. and CHATTERJEE, A.K. Extracellular enzymes and pathogenesis of soft rot *Erwinia*. *Annual Review of Phytopathology*, September 1994, vol. 32, p. 201-234.

BEVAN, M. Binary vectors for plant transformation. *Nucleic Acids Research*, November 1984, vol. 12, no. 22, p. 8711-8721.

BRADFORD, M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, May 1976, vol. 72, p. 248-254.

BREWBAKER, J.L. and KWACK, B.H. The essential role of calcium ion in pollen germination and pollen tube growth. *American Journal of Botany*, 1963, vol. 50, p. 859-865.

BRUCE, W.B.; DENG, X.W and QUAIL, P.H. A negatively acting DNA sequence element mediates phytochrome-directed repression of *phyA* gene transcription. *EMBO Journal*, October 1991, vol. 10, no. 10, p. 3015-3024.

CANEVASCINI, S.; CADERAS, D.; MANDEL, T.; FLEMING, A.J.; DUPUIS, I. and KUHLEMEIER, C. Tissue-specific expression and promoter analysis of the tobacco *ltp1* gene. *Plant Physiology*, October 1996, vol. 112, no. 2, p. 513-524.

CHAMBERLAND, H.; OUELLETTE, G.B.; PAUZE, F.J. and CHAREST, P.M. Immunocytochemical localization of tomato pectinesterase in root cells of tomato plants infected by *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Canadian Journal of Botany*, June 1991, vol. 69, no. 6, p. 1265-1274.

CHEN, M.H. and CITOVSKEY, V. Systemic movement of tobamovirus requires host cell pectin methylesterase. *Plant Journal*, August 2003, vol. 35, no. 3, p. 386-392.


CHRISTGAU, S.; KOFOD, L.V.; HALKIER, T.; ANDERSEN, L.N.; HOCKAUF, M.; DORREICH, K.; DALBOGE, H. and KAUPPINEN, S. Pectin methyl esterase from *Aspergillus aculeatus*: expression, cloning in yeast and characterization of the recombinant enzyme.

- Biochemical Journal*, November 1996, vol. 319, no. 3, p. 705-712.
- COLLMER, A. and KEEN, N.T. The role of pectic enzymes in plant pathogenesis. *Annual Review of Phytopathology*, September 1986, vol. 24, p. 383-409.
- DARASELIA, N.D.; TARCHEVSKAYA, S. and NARITA, J.O. The promoter for tomato 3-hydroxy-3-methyl-glutaryl coenzyme a reductase gene 2 has unusual regulatory elements that directs high-level expression. *Plant Physiology*, October 1996, vol. 112, no. 2, p. 727-733.
- DITTA, G. A new broad-host range DNA cloning vector for use with *Rhizobium* and other gram-negative bacteria. In: PANOPOULOS, N.J. ed. *Genetic Engineering in the Plant Sciences*. New York, USA. Praeger Publishers. 1981, p. 145-154.
- DUWE, B. and KHANH, N.Q. Site-directed mutagenesis of the active site of pectin methylesterase from *Aspergillus niger* RH5344. *Biotechnology Letters*, June 1996, vol. 18, no. 6, p. 621-626.
- FELSENSTEIN, J. PHYLIP (Phylogeny Inference Package) version 3.5c. *Department of Genetics, University of Washington, Seattle*, 1993.
- FELSENSTEIN, J. PHYLIP - Phylogeny Inference Package (Version 3.2). *Cladistics*, March 1989, vol. 5, no. 1, p. 164-166.
- FLUHR, H.J. and DANDEKAR, A.M. Organ specific and light-induced expression of plant genes. *Science*, May 1986, vol. 232, no. 4754, p. 1106-1111.
- FRENKEL, C.; PETERS, J.S.; TIEMAN, D.M.; TIZNADO, M.E. and HANDA, A.K. Pectin methylesterase regulates methanol and ethanol accumulation in ripening tomato (*Lycopersicon esculentum*) Fruit. *Journal of Biological Chemistry*, February 1998, vol. 273, no. 8, p. 4293-4295.
- GAFFE, J.; TIZNADO, M.E. and HANDA, A.K. Characterization and functional expression of a ubiquitously expressed tomato pectin methylesterase. *Plant Physiology*, August 1997, vol. 114, no. 4, p. 1547-1556.
- GAFFE, J.; TIZNADO, M.E. and HANDA, A.K. Cloning and nucleotide sequence of a pectin methylesterase cDNA homologue (Accession No. U49330) from tomato leaves (PGR96-017). *Plant Physiology*, April 1996, vol. 110, no. 4, p. 1436.
- GAFFE, J.; TIZNADO, M.E. and HANDA, A.K. Cloning and characterization of a pectin-methylesterase gene expressed throughout growth and development of tomato plants. In: *Annual Meeting of the American Society of Plant Physiologists*. (29<sup>th</sup> July-2<sup>nd</sup> August, 1995, Charlotte, North Carolina, USA). Plant Physiology, Rockville, MD, USA. p. 76.
- GREEN, P.J.; KAY, S.A. and CHUA, N.H. Sequence-specific interaction of a pea nuclear factor with light-responsive elements upstream of the rbcS-3A gene. *EMBO Journal*, September 1987, vol. 6, no. 9, p. 2543-2549.
- GUILTINAN, M.J.; MARCOTTE, Jr., W.R. and QUATRANO, R.S. A plant leucine zipper protein that recognizes an abscisic acid response element. *Science*, October 1990, vol. 250, no. 4978, p. 267-271.
- HARRIMAN, R.W.; TIEMAN, D.M. and HANDA, A.K. Molecular cloning of tomato pectin methylesterase gene and its expression in rutgers, ripening inhibitor, nonripening, and never ripe tomato fruits. *Plant Physiology*, September 1992, vol. 97, no. 1, p. 80-97.
- HOOYKAAS, P.J.J. Transformation of plant cells via *Agrobacterium*. *Plant Molecular Biology*, September 1989, vol. 13, no. 3, p. 327-336.
- JEFFERSON, R.A.; KAVANAGH, T.A. and BEVAN, M.W. GUS Fusions: b-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO Journal*, December 1987, vol. 6, no. 13, p. 3901-3907.
- JENKINS, J.; MAYANS, O.; SMITH, D.; WORBOYS, K. and PICKERSGILL, R.W. Three-dimensional structure of *Erwinia chrysanthemi* pectin methylesterase reveals a novel esterase active site. *Journal of Molecular Biology*, January 2001, vol. 305, no. 4, p. 951-960.
- JOSHI, C.P. An inspection of the domain between putative TATA box and translational start site in 79 plant genes. *Nucleic Acid Research*, August 1987, vol. 15, no. 16, p. 6643-6648.
- KIM, H.U.; PARK, B.S.; JIN, Y.M. and CHUNG, T.Y. Promoter sequences of two homologous pectin esterase genes from chinese cabbage (*Brassica campestris* L. ssp. Pekinensis) and pollen-specific expression of the GUS gene driven by a promoter in tobacco plants. *Molecules and Cells*, February 1997, vol. 7, no. 1, p. 21-27.
- KO, C.H.; BRENDEL, V.; TAYLOR, R.D. and WALBOT, V. U-Richness is a defining feature of plant introns and may function as an intron recognition signal in maize. *Plant Molecular Biology*, March 1998, vol. 42, no. 4, p. 573-583.
- LABEL, P.; IMBAULT, N. and VILLAR, M.E. Quantitation and GC-MS identification of abscisic acid in stigma, ovary and pedicel of pollinated poplar flowers (*Populus nigra* L.). *Tree Physiology*, May 1994, vol. 14, no. 5, p. 521-530.
- LI, Q. and HUNT, A.G. A near-upstream element in a plant polyadenylation signal consists of more than six

- nucleotides. *Plant Molecular Biology*, August 1995, vol. 28, no. 5, p. 927-934.
- LIU, H.X. and FILIPOWICZ, W. Mapping of branchpoint nucleotides in mutant mRNAs expressed in plant cells. *Plant Journal*, January 1996, vol. 9, no. 2, p. 381-389.
- LINEWEAVER, H. and JANSEN, E.F. Pectic Enzymes. In: NORD, F.F. ed. *Advances in Enzymology*, Volume XI, New York, Interscience Publishers, Inc., 1951, p. 100-150.
- MA, R.; REESE, J.C. BLACK, W.C. IV and BRAMEL-COX, P. Detection of pectinesterase and polygalacturonase from salivary secretions of living greenbugs, *Schizaphis graminum* (Homoptera:Aphididae). *Journal of Insect Physiology*, March 1990, vol. 36, no. 7, p. 507-512.
- MANDEL, T.A.; FLEMING, A.J.; KRAHENBUHL, R. and KUHLEMEIER, C. Definition of constitutive gene expression in plants: the translation initiation factor 4a gene as a model. *Plant Molecular Biology*, December 1995, vol. 29, no. 5, p. 995-1004.
- MATHIS, N.L. and HINCHEE, M.A.W. *Agrobacterium* Inoculation technique for plant tissues. In: GELVIN, S.B. and SCHILPEROORT, R.A. eds. *Plant Molecular Biology Manual*, Dordrecht, Kluwer Academic Publishers, 1994, p. B6:1-9.
- MENDGEN, K.; HAHN, M. and DEISING, H. Morphogenesis and mechanisms of penetration by plant pathogenic fungi. *Annual Review of Phytopathology*, September 1996, vol. 34, p. 367-386.
- MU, J.H.; STAINS, J.P. and KAO, T.H.. Characterization of a pollen-expressed gene encoding a putative pectin esterase of *Petunia inflata*. *Plant Molecular Biology*, June 1994, vol. 25, no. 3, p. 539-544.
- MYERS, E. and MILLER, W. Optimal alignments in linear space. *Bioinformatics*, March 1988, vol. 4, no. 1, p 11-17.
- NAIRN, C.J.; LEWANDOWSKI, D.J. and BURNS, J.K. Genetics and expression of two pectinesterase genes in valencia orange. *Physiologia Plantarum*, February 1998, vol. 102, no. 2, p. 226-235.
- NAP, J.P.; KEIZER, P. and JANSEN, R. First-generation transgenic plants and statistics. *Plant Molecular Biology Reporter*, June 1993, vol. 11, no. 2, p. 156-164.
- NESTER, E.W.; KEMNER, J.; DENG, W.; LEE, Y.W.; FULLNER, K.; LIANG, X.; PAN, S. and HEATH, J.D. *Agrobacterium*: A natural genetic engineering exploited for plant biotechnology. In: STACEY, G., MULLIN, B. and GRESSHOFF, P.M. eds. *Biology of Plant-Microbe Interaction*. St. Paul, Minnesota, 1996, p. 111-119.
- PLASTOW, G.S. Molecular cloning and nucleotide sequence of the pectin methyl esterase gene of *Erwinia chrysanthemi* B374. *Molecular Microbiology*, March 1988, vol. 2, no. 2, p. 247-254.
- PRESTIDGE, D.S. Signal SCAN: A computer program that scans DNA sequences for eukaryotic transcriptional elements. *Bioinformatics*, April 1991, vol. 7, no. 2, p. 203-206.
- QUANDT, K.; FRENCH, K.; KARAS, H.; WINGENDER, E. and WERNER, T. MatInd and MatInspector - New fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acid Research*, December 1995, vol. 23, no. 23, p. 4878-4884.
- REXOVA-BENKOVA, L. and MARKOVIC, O. Pectic enzymes. In: TIPSON, R.S. and HORTON, D. eds. *Advances in Carbohydrate Chemistry and Biochemistry*, Academic Press, Inc. Massachusetts, 1976, p. 323-385.
- RICARD, J. and NOAT, G. Electrostatic effects and the dynamics of enzyme reactions at the surface of plant cells. 1. A theory of the ionic control of a complex multi-enzyme system. *European Journal of Biochemistry*, February 1986, vol. 155, no. 1, p. 183-190.
- RICHARD, L.; QIN, L.X. and GOLDBERG, R. Clustered genes within the genome of *Arabidopsis thaliana* encoding pectin methylesterase-like enzymes. *Gene*, May 1996, vol. 170, no. 2, p. 207-211.
- ROGER, D.; LACOUX, J.; LAMBLIN, F.; GAILLET, D.; DAUCHEL, H.; KLEIN, D.; BALANGE, A.P.; DAVID, A. and LAINE, E. Isolation of a flax pectin methylesterase promoter and its expression in transgenic tobacco. *Plant Science*, March 2001, vol. 160, no. 4, p. 713-721.
- ROYO, J.; DIAZ, I.; RODRIGUEZ-PALENZUELA, P. and CARBONERO, P. Isolation and promoter characterization of barley gene *Itr1* encoding trypsin inhibitor BTI-CMe:differential activity in wild type and mutant *lys3a* endosperm. *Plant Molecular Biology*, 1996, vol. 31, p. 1051-1059.
- SAMBROOK, J.; MANIATIS, T. and FRITSCH, E.F. *Molecular Cloning: A Laboratory Manual*. New York, Cold Spring Harbor Laboratory Press, 1989.
- SCHLINDER, U.; MENKENS, A.E.; BECKMAN, H.; ECKER, J.R. and CASHMORE, A.R. Heterodimerization between light-regulated and ubiquitously expressed arabidopsis GBP vZIP Proteins. *EMBO Journal*, April, 1992, vol. 11, no. 4 , p. 1261-1273.
- SEXTON, R. and ROBERTS, J.A. Cell biology of abscission. *Annual Review of Plant Physiology*, June 1982, vol. 33, p. 133-162.
- SHEN, Z.C.; MANNING, G.; REESE, J.C. and REECK, G.R. Pectin Methylesterase from the Rice Weevil, *Sitophilus oryzae* (L.) (Coleoptera:Curculionidea):

- Purification and characterization. *Insect Biochemistry and Molecular Biology*, March 1999, vol. 29, no. 3, p. 209-214.
- STEPHENSON, M.B. and HAWES, M.C. Correlation of pectin methylesterase activity in root caps of pea with root border cell separation. *Plant Physiology*, October 1994, vol. 106, no. 2, p. 739-745.
- THOMPSON, J.D.; HIGGINS, D.G.; GIBSON, T.J. and CLUSTAL. W. Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, November 1994, vol. 22, no. 22, p. 4673-4680.
- TIEMAN, D.M. and HANDA, A.K. Reduction in pectin methylesterase activity modifies tissue integrity and cation levels in ripening tomato (*Lycopersicon esculentum* mill.) Fruits. *Plant Physiology*, October 1994, vol. 106, no. 2, p. 429-436.
- TURNER, L.A.; HARRIMAN, R.W. and HANDA, A.K. Isolation and nucleotide sequence of three tandemly arranged pectin methylesterase genes (accession Nos. U70675, U70676 and U70677) from Tomato. *Plant Physiology*, November 1996, vol. 112, no. 3, p. 1398.
- VALETTE-COLLET, O.; CIMERMAN, A.; REIGNAULT, P.; LEVIS, C. and BOCCARA, M. Disruption of *Botrytis cinerea* pectin methylesterase gene Bcpme1 reduces virulence on several host plants. *Molecular Plant-Microbe Interactions*, April 2003, vol. 16, no. 4, p. 360-367.
- WIETHOLTER, N.; GRAESSNER, B.; MIERAU, M.; MORT, A.J. and MOERSCHBACHER, B.M. Differences in the methyl ester distribution of homogalacturonans from near isogenic wheat lines resistant and susceptible to the wheat stem rust fungus. *Molecular Plant-Microbe Interactions*, October 2003, vol. 16, no. 10, p. 945-952.
- WILSON, C.; BELLEN, H.J. and GEHRING, W.J. Position effects on eukaryotic gene expression. *Annual Review of Cell Biology*, November 1990, vol. 6, p. 679-714.
- WINGENDER, E.; KEL, A.E.; KEL, O.V.; KARAS, H.; HEINEMEYER, T.; DIETZE, P.; KN'UPPEL, R.; ROMASCHENKO, A.G. and KOLCHANOV, N.A. TRANSFAC, TRRD and COMPEL: Towards a federated database system on transcriptional responsive element. *Nucleic Acids Research*, January 1997, vol. 25, no. 1, p. 265-268.
- WINGENDER, E.; DIETZE, P.; KARAS, H. and KN'UPPEL, R. TRANSFAC: A database on transcription factors and their dna binding sites. *Nucleic Acids Research*, January 1996, vol. 24, no. 1, p. 238-241.
- WEN, F.; ZHU, Y. and HAWES, M.C. Effect of pectin methylesterase gene expression on pea root development. *Plant Cell*, June 1999, vol. 11, no. 6, p. 1129-1140.
- ZENG, Y.R.; PANDEY, M.; PRASAD, N.K. and SRIVASTAVA, G.C. Hydrolysing enzymes and respiration during ripening of tomato (*Lycopersicon esculentum*) fruits. *Current Science*, June 1996, vol. 70, no. 11, p. 1017-1019.
- ZHU, J.K. Plant salt tolerance. *Trends in Plant Science*, February 2001, vol. 6, no. 2, p. 66-71.
- ZUPAN, J.R. and ZAMBRYSKI, P. Transfer of the T-DNA from *Agrobacterium* to the plant cell. *Plant Physiology*, April 1995, vol. 107, no. 4, p. 1041-1047.

## APPENDIX Figures



**Figure 1. Partial restriction map of the  $\lambda$  phage and location of the PMEU1 genomic sequence.** Open box, black box and gray box represent the PMEU1 promoter, genomic DNA coding region, and the phage DNA region flanking the 3' end of *pmeu1* gene. Right and left represent the right (8.8 kb) and left (19.9 kb) lambda arms. Abbreviations: B, E, H,P, and S indicate BamHI, EcoR I, Hind III, Pst I and Sal I restriction sites. Sal I sites at the left and right borders are from the  $\lambda$  EMBL3A.

-79 GGACCAATGT CACGGATATA AAACCCCCAC CAATCCGATC CAATTTCTCC  
-31 ACAACTCTCC CTTAAATTTT TCCATCCAAA ATGACACGTG TTGAAGATTT  
21 TTTCAGCAAA CAAATCGATT TTTGTAAAAG GAAGAAAAAA ATCTACTTGG  
71 CCATTGTTGC CTCAGTCTTG CTGGTTGCTG CAGTAATCGG AGTAGTCGCC  
121 GGAGTAAAAT CTCATTGAA AACTCCGAC GATCATGCAG ACATAATGGC  
171 CATTTCGTCT TCAGCCCATG CTATTGTAAA ATCTGCGTGT AGCAACACTC  
221 TACACCCCGA ACTGTGTAC TCTGCGATTG TCAATGTTTC TGATTTCTCA  
271 AAAAAAGTAA CAAGCCAAAA AGATGTGATT GAATTGTCTT TGAATATCAC  
321 TGTCAAAGCC GTTCGACGCA ACTACTATGC AGTCAAGGAA CTCATCAAAA  
371 CTAGAAAAGG TACTTGRCRC GTTARCTTA ATACTCCATT GTABBBATGT  
421 TBSAAGTICT CTTCTTCTTT TATTGAAAT ATAGTTGGTT TAICTCACTA  
471 TTTTATTTTT CATAGGTTTA ACCCCACGAG AAAAGGTTGC GCTGCATGAC  
521 TCCTTGGAGA CGATGGACGA GACACTCGAC GAGCTCCACA CTGCTGTAGA  
571 AGATCTGGAG CTATATCCCA ACAAAAAATC ATTGAAAGAA CACGTGCAAG  
621 ACCGTAAAAC TCTAATAAGT TCCGCAATTA CAAACCAGGA AACTTCCGCTC  
671 GACGGTTTTCT CTCACGATGA GCGCGATAAA AAGGTACGCA AGTTTTTGT  
721 GAAAGGCCAA AAGCACGTGG AAAAAATGTG CAGCAATGCT TTAGCTATGA  
771 TCTGTAACAT GACCGATACC GACATTGCAA ATGAGATGAA ATTATCGGCC  
821 CCCGCCAATA ATAGGAAGTT AGTAGAGGAT AACGGCGAGT GCGCGAGTG  
871 GTTGTCCGCC GCGACAGGA GGTATTGCA GTCGTCGACG GTGACGCGAG  
921 ATGTGGTTGT GCGCGCCGAC GGAAGCGGAG ATTACAAAAAC GGTCGACAG  
971 GCGGTACGAA AAGCGCCAGA GAAGAGTAGC AAGAGGTATG TGATTAGGAT  
1021 AAAAGCTGGT GTTACAGGG AAAACGTGGA TGTGCCAAAG AAGAAGACGA  
1071 ATATTATGTT TATGGGAGAT GCAAAAAGCA ATACAATAAT CACAGCAAGT  
1121 AGGAATGTGC AAGATGTTAG CACTACCTTC CACTCTGCTA CAGTTGGTBA  
1171 GTTATTATTA TTATCTTTAT CAACCAATTG CCTTAATTG CAGCTAECTA  
1221 CTTATACBAG GTAGAGTTAA ATTTAATTTG GTAGCGAGT GATBTAATTT  
1271 TGTATCACAT GTTAAATGTA TACTAATTTT TTACTTTANT ACTTTATAGA  
1321 TAGTCAAGGA ACAGTAAAG TGAAACCAAT AACACACTTT CATTCCGCTG  
1371 CTAGATAGT GARGCAATA AACACACTTC ACTTAATAT TTGTCAAGT  
1421 GGTAGCATT TAGGTTGATC TATTCTCCTT GTTAAATAAA AGAAGTTCAC  
1471 TATTTACCT TTAAAAGGAG AAGGATTAAT TAAATTAGAC TTGACCAAG  
1521 AGTTCGAGTA CTAAATAGT TCAGTAATAG ACTATTCTTG TTGGTAGTA  
1571 GTTAGAGGA ATTAAGGAGC TTTTGTACT ACAATAGTC GTATAATTAA  
1621 AGTAAAGTT TATCTCTTA TTGTACTACA AATAGAGAGA AATAATTTCA  
1671 CTTATACCA AAAACATAT ACATACTCTT GTTTTTTTAA TCCTGAGATT  
1721 ATTATCCCTT AACCAATCAA CCAACAGTT AATGTTATTG GTGAAGGCTG  
1771 GCTTATACTC AATATGACTA TATATAACGA ATCCACGTC TGTAAGGTT  
1821 GGATAAAGG CATATCTTAT GGACBAGG ATATAGATTI GTCBCCACT  
1871 TTTTGTAGT GGTGTATCTA TCBGCTTGTG ATATAGATC TGCTTTCCBG  
1921 CTGGGGTAGC GCTCCCTTTC CFTTGAATAA TTCATTAATG ACTTCTTCT  
1971 TGTATAATA TTTTTACTA CATATAATTT TCGTACCBA GGTCCGTTTT  
2021 CTTTGCAGCG GCCTATAATA ATCGTGTCTT TTGATAAGGA CAGCTCAGGG  
2071 GTACATTCTT AGCTGACRAAT AATAGAGACA CCCCCTCTT TAGTCCGCTGA  
2121 GATATTTATT CAARRATTA AATGTATCA TGACRAATRT TANTGAGTTG  
2171 TCCAAATAG AATGACATA ACTAATTTGA TCGBTCCGC GTGGCAGGAA  
2221 AAGTTCTTGC CCGGATATA ACCTTCCAAA ACACAGCAGG AGCCTCGAAG  
2271 CATCAAGCCG TGCACTCTG CGTGGGCTCT GATTTGTCCG CATTTHATAG  
2321 ATGTGACATG TFGGCTTATC AGGACACCCT CTACGTCCAC TCTAATCGTC  
2371 AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA

```

AtPME9      1 -----
AtPME10     1 -----
AtPME8      1 -----
ATPME4      1 -----MIGKVVVSVASILLVGVVAIGVVAFINKN
BP19        1 -----MAVGKIVISVASMLLVGVVAIGVVTFVNKG
PER          1 -----MGGTRYNGGHDOSKRFALVGVSSILLVAVVATVADAQQ--
PEF1        1 -----MGGNDNNGGQGGQKKHALLVGSCILLVAVVGVVAVSLTKG
PPE1        1 -----
AtPME6      1 -----MGSDGDKKKKIVAGSVSGFIVIMVSVAVVTSKH
ATPME2      1 -----EFISKFSDFKNNKKLILSSAAIALLLLASTVIGIAATTTNQNKNOK
ATPME3      1 -----MAPSMKEIFSKDNFKNNKKLVLLSAAVALFVAAVAGISAGASKANEKR-
PECS-1.1    1 -MTHIKEFFTKLSESSSNQINISNIPKKKKLFLALFATLLVVAAVVIGVAGVNSRKNSGD
PECS-1.2    1 -----
PMEU1       1 -----MTRVEDFFSKQIDFCRKKKKIYLAIVASVLLVAAVVIGVVAGVKSHSKNSD
PECS-2.1    1 -----MALRILITVSLVLFSLSHTSPGYSP
LEPME1      1 -----
LEPME2      1 -----MATPQQPLLTKHKQNSIISFKILTFVVTIFVALFLVVFLVAPY
LEPME3      1 -----MATPLQPFLTRKHKQNP IIGFNILTFVVTIFVALFLVVFLVAPY
RCPME1      1 -----MAIQETLIDPKKSPKPTFWLLLSLAAVIGSSALIVSHLNKPI
AtPME7      1 -----
OsPME       1 MAHATLGSPEPAAPRLCADGRHRRRLIVVLCIVGVAVAVAVAVLGRSRMTSS
ATPME1      1 ---MDSVNSFKGYGKVDEAQLALKKKTRKRLLLSISVVVLIAMVIAAVVAVTVVHKVKN
PMEB        1 -----

AtPME9      1 -----
AtPME10     1 -----
AtPME8      1 -----
ATPME4      30 GDAN---LSPQMKAVQGIQSTSDKASCVKTEPVKSE---DPNKLKAFMLATKDE
BP19        31 GGAGGDKTLNSHQKAVESLCASATDKGSCAKTIDPVKSD---DPSKLKAFMLATKDA
PER          39 -----GQPNVQILCESTQYQQTCHQSLAKAPAETAGVKDLIKAAFSAT---SE
PEF1        41 GDGEQKAHISNSQKNVDMLCQSTKFKETCHKTEKASFS--NMKNRIKGAIGAT---EE
PPE1        1 -----
AtPME6      36 SPRDDENHIRKTTKAVQAVCAPTDFKDTCVNSIMGASPDSDDPVDLIKLGFKVT---IK
ATPME2      46 ----ITTLFSTSHAILKSVCSSTLYPELCFSAVAATGGK--ELTSOKEVIEASLNLTTKA
ATPME3      50 -----TLPSSHAVLRSSCSSTRYPCLCISAVVTAGAC--ELTSQKDVIEASVNLTTITA
PECS-1.1    60 -----NGNEPHHAILKSSCSSTRYPDLCFSAVAAVPEASKKVTSQKDVIEMSLNITTTA
PECS-1.2    1 -----
PMEU1       51 DHADIMAISSSAHAIDKSAACSNLHPELCVSAIVNVVSDFSKKVTSQKDVIELSLNITVKA
PECS-2.1    27 -----EVKSWCGKTPNPQPCFYFLTQKTDVT---SIKQDTEFYKISLQLAL
LEPME1      1 -----
LEPME2      45 Q-----FEIKHSN--LCKTAQDSQLCLSYVSDLMS-NEIVTTSDGSLILMKFLVNY
LEPME3      45 Q-----FEIKHSN--LCKTAQDSQLCLSYVS-----EIVTTSDGVTVLKFLVKY
RCPME1      44 S-----FFPLSSAPNLCEHAVDTKSLCTHSEVVOGQALANTRDKHLSTLISLLTKS
AtPME7      1 -----
OsPME       61 SGGGRAPRGRAPTEAARTCGVTLYPELCVGEIMAFPGAAG--AGDAELVPMSLNATHRR
ATPME1      58 ESTPSPPELTPSTSKAICSVTRFPESCISSESKLPSS---NTTDPETLFLKLSLKI
PMEB        1 -----

AtPME9      1 -----
AtPME10     1 -----
AtPME8      1 -----
ATPME4      81 LTKSSNFTGQTEVNMGSSISPNNKAVLDYCKRVFMYALDDIATIIEEMG---EDL
BP19        86 VTKSTNFTASTEEMGRNINATSKAVLDYCKRVLMYALDDIETIIEEMG---EDL
PER          84 ELLKHINSS--LIQELCQDKMTKQAEVQNEVLDYAVDGHKSGAVD---KFDI
PEF1        95 ELRKHINNSA--LYOELATDSMTKQAEICNEVLDYAVDGHKSGVGLD---QDFD
PPE1        1 -----
AtPME6      92 SINESLEKASGDIKAKADKNPEAKGAFELCEKLMIDAIDDKCKDHDG-----FSV
ATPME2      100 VKHN-YFAVKKLIAKRGLTPREVTALHDCLDTEIDETLDELHVAVEDLHQY---PKQ-
ATPME3      102 VEHN-YFTVKKLIKRRGLTPREKTAALHDCLDTEIDETLDELHETVEDLHLY---PTK-
PECS-1.1    114 VEHN-YFGIQKLLKRTN-LTKREKVALHDCLDTEIDETLDELHKAQEDLEEY---PNK-
PECS-1.2    1 -----
PMEU1       111 VRRN-YYAVKELIKTRKGLTPREKVALHDCLDTEIDETLDELHTAVEDLELY---PNK-
PECS-2.1    70 ERAT-TAQSRTYTLGSKCRNEDEKAVEDCRELYELTVLKNQTSN-----
LEPME1      1 -----NDIRQHGALTDCLLDDQSVDLASDSIAATD-----
LEPME2      94 VHQMNAIPVVKMKMNQINDIQEGALTDCLLDDQSVDLASDSIAATD-----
LEPME3      89 VHQMNAIPVVRKIKNQINDIQOQALTDCLLDDQSVDLASDSIAATD-----
RCPME1      96 TSHIQKAMETANVIKRRVNSPKETALNDCQLMDLSEDRVWDSQLTET-----
AtPME7      1 -----MKERKCFADPFGDRRLQNAVSDCLLDFSSBETWSASASENP-----KGGK
OsPME       119 VVDALYNATALGGAALLAGARSGAAYGDCVEMLDAAEELIARSUGATAAPPPPDSDVA
ATPME1      113 IDELDSISDLPEKLSKETEDERIKSALRVCGDLIEDALDRNDTWSAIDDE----EKKK
PMEB        1 -----MSLTHYSGLAAAAMSLLITACGGQTPNSARFQPVFPGTVSRP-----VL

```

Isolation and study of a ubiquitously expressed tomato pectin methylesterase regulatory region

AtPME9 1 ---MGYISLALVALLVFFASPVVLAADDITPIPADRAQIPQWFMANVFPFSQRRGTDPPEL  
 AtPME10 1 ---MGYISMSVVAFLVVFASPVVLAATDIPENRAQIPQWFKTNVFPYSQRRGTDPAL  
 AtPME8 1 -----MVFAADDLTP IPEGKPVVQVFNTHVGPLAQKRGDPAL  
 ATPME4 133 SQIGSKIDQLKQNLIGVYNYQTDCLDDIEEDDLRKAIGEGIANSKILTTNAIDIFHTVVS  
 BP19 138 QQSGSKMDQLKQNLIGVYNYQTDCLDDIEESELRKVMGEGIAHSKILSSNAIDIFHALTT  
 PER 135 NKIHEYSYDLKRWLIGLSSHQQTCLDGFANTTTKAGETMARALNTS IQLSSNAIDVVDVAV  
 PEF1 146 HKLSEYAFDIKRWLIGLSSHQQTCLDGFVNTKTHAGETMAKVLKTSMESSNAIDVMDVV  
 PPE1 1 -----MVKLLNSTFELSINALSMLNSF  
 AtPME6 143 DQIEVFVEDLRYWLSGSIAFQQTCDMSFGEIKSNLMQDMLKIFKTSSELSNSLAVVTRI  
 ATPME2 153 KSLRKHADDLKTLISSAINTQQTCLDGFSDADRVRKALLKGOVVEVHMCSNAIAMIK  
 ATPME3 155 KTLREHAGDLKTLISSAINTQQTCLDGFSDADKQVRKALLKGOVVEVHMCSNAIAMIK  
 PECS-1.1 166 KLSLQHADDLKTLMSSAANTNQQTCLDGFSDADANKHVRDALSDGOVVEVKMCSNAIAMIK  
 PECS-1.2 1 -----IK  
 PMEU1 164 KSLKEHVEDLKTLLISSAINTQQTCLDGFSDADKVRKALLKGOVVEVKMCSNAIAMIK  
 PECS-2.1 115 SSPGCTKVDKQITLSTALTNLETCRASLEDLGVPEYVLPPLSN-----AKVALAMLASITTT  
 LEPME1 32 KRSRSEHAMSESWLSGVLTNHVTCIDELDSFTKAMINGTNLEELISEAKVALAMLASITTT  
 LEPME2 143 KRTHSEHAMASWLSGVLTNHVTCIDELDSFTKAMINGTNLEELISEAKVALAMLASVTT  
 LEPME3 138 KRSRSEHAMASWLSGVLTNHVTCIDELTSFSLSTRNGTVLDELITSEAKVALAMLASVTT  
 RCPME1 145 KMNIDSQDAHTWLSGVLTNHATCLNGLEGTSRVVMESSDQDLISEARSSLAVALSVLP  
 AtPME7 50 NGTGDVGSDFRITWLSAALSNOQTCEMGFDGTSGLVKSLSVAGSLDQLYSMLRELLPVPQPE  
 OsPME 179 DTAGRDDDDIMTWLSAALSNSHDTCEMDSLQEVGAGGDDGDDGGRIKQMLGYLGNLGEHL  
 ATPME1 168 TLSSSKIEDLKTWLSAALVTDHETCFDLSDELKONKTEYANSTITQNLKMSMRSTFTSN  
 PMEB 46 SAQEAGRFTPOHFFAHGGEYAKPVADGWTPPTIDTSRVTAAYVVGPRAGVAGATHTSIQQ

AtPME9 58 EAAEAS-----  
 AtPME10 58 EAAEAA-----  
 AtPME8 39 VAAEAA-----  
 ATPME4 193 AMAKINNKVDDLKMNMTGGIPTPGAPPVVDESVPADPDGPARRLLEDIDETGIPITWVSGAD  
 BP19 198 AMSQMNKVVDDMK-----KGNLGETPAPDRDLLEDLDQKGLPKMHSDDKD  
 PER 195 YDLTN-----AKRLLSLDNG-----YPLWVSEGQ  
 PEF1 206 SRILKGFHPSQYGVSRLLSDG-----IPSWVSDGH  
 PPE1 23 GDMVAQATG---LNRKLLTTDS-----SDATA  
 AtPME6 203 STLIPNSNLTAKYARKLLSTEDS-----IPTWVSGPEA  
 ATPME2 213 NMTETDIANFELRDKFFNLHQQQQRK-----LKEVTG-----DLSDGVPKMLSVGD  
 ATPME3 215 NMTDTDIANFEQKAKITSNRRLKEENQETTVAVDIAGAG-----ELDSEGEPTWVSGD  
 PECS-1.1 226 NMTDTDMMIMRTSNRRLTEETSTVDG-----WPAWVSPGD  
 PECS-1.2 3 NMTDTDMMIMRTSNRRLIEETSTVDG-----WPAWVSTGD  
 PMEU1 224 NMTDTDIANEMKLSAPANNRKLVEDNGE-----WPEWVSGD  
 PECS-2.1 158 -NVTKLISNTLSLNKVPYNEPSYKDG-----FETWVSPGD  
 LEPME1 92 QDED---VFMTGLGKMP-----SWVSSMD  
 LEPME2 203 PNDD---VLRPGLGKMP-----SWVSSRD  
 LEPME3 198 PNDE---VLRQGLGKMP-----YVWSSRD  
 RCPME1 204 AKSMDGFIDESLNGEFP-----SWVTSKD  
 AtPME7 110 QKPKAVSKPGPIAKGPKAPPGRKLRDTE-----EDESLOFPDWRPDD  
 OsPME 239 SNSLAIFAARGRPGGELSDVPVHNQLHR-----RLLTIDDDDDDDGSPFRWRHMD  
 ATPME1 228 SLAIVSKILSALSDLGPIIHRRRRLMS-----HHHQQSVDFEKWAR  
 PMEB 106 AVNAALRQHPGQTR-----VYIKLLP

AtPME9 64 -----RRVITVNVQNGGDFKTNAAIKSIFLANKNEV  
 AtPME10 64 -----RQIITVNVQNGGDFKTNAAIKSIFLANKNEV  
 AtPME8 45 -----PRIINWNPKGG-EFKTLTDATKSVFAGNTRRV  
 ATPME4 253 RRLMAKAGR-----GRRGGRRGGARVRTNFVVAKDGGGQFKTVQQAADACPENNRGRC  
 BP19 242 RRLMAQAGRPGAPADEGIGEGGGGGKIKPTEVVAKDGGGQFKTISEAVKAKCEKKNPGRC  
 PER 220 RRLLAEAT-----VKPNVVAQDGGGQFKTLTDATKTVFANNAQNF  
 PEF1 238 RRLLAGGN-----VKNANVVAQDGGGQFKTLTDALKTVPPTNAAPF  
 PPE1 47 RRLQIISN-----AKPNATVVAQDGGGQFKTIKALDAVPEKKNTEPF  
 AtPME6 235 RRLMAAQGGG-----PGPVKANAVVAQDGTGQFKTITDALNAVPEKGNKVPF  
 ATPME2 260 RRLQGS-----TIKADATVADGGGDFDNGSAVAAAPKSNRRF  
 ATPME3 270 RRLQGS-----GVRDATVADGGGDFKTVAAVAAAPENSRRY  
 PECS-1.1 262 RRLQSS-----SWTPMAVVAADGGGDFKTVAAVAAAPQGGTRRY  
 PECS-1.2 39 RRLQSS-----SWTPMVVAADGGGDFKTVAAVAAAPQGGTRRY  
 PMEU1 261 RRLQSS-----TWTPDVVAADGGGDFKTVSEAVRKAPEKSSRRY  
 PECS-2.1 192 RRLQQT-----TPRANVVAQDGGGDFKTVAAVAAAPSRAGGSRY  
 LEPME1 113 RRLMESSGK-----DIIANRVVAQDGTGKYFTLAEAVAAAPNRSKRY  
 LEPME2 224 RRLMESSGK-----DIIANRVVAQDGTGKYFTLAEAVAAAPDKNKTRY  
 LEPME3 219 RRLMESSGK-----DIIANRVVAQDGTGDTCTLAEAVAAAPDKNKTRY  
 RCPME1 228 RRLLESTVG-----DIKANVVAQDGGGDFKTVAAVAAAPDNGKARY  
 AtPME7 153 RRLLESN-----GRTYDVSVALDGTGNTKIMDAIKKAPDYSSTRF  
 OsPME 290 RRLQAAAA-----ETIADMVVAQDGTGTHKIRDAIKKAPHSRRV  
 ATPME1 269 RRLQQTAG-----LKPDTVVAQDGTGDLTVNEAVAKVKKSLKMF  
 PMEB 127 GTYTGTVYVP-----EGAPPITLFGAGDRPEQVVSIALDLSMMSPADY

```

AtPME9      96  IIKLAPGTYHERKVTVDVGRPYVTLLEGKPGAETMLTYGTAAKYG---TVESATLIWVWATN
AtPME10     96  IIKLAPGVYNEKVTIDLARPFITLLGQPGAETVLYHGTAAQYG---TVESATLIWVWEY
AtPME8      76  IIKMAHCEYREKVTIDEMKPFITLHGQPNAMPVITYDCTAAKYG---TVDSASLIILSDY
ATPME4      306 IYIKAGLYREQVITPRKKNNVFMFGDGRKTVISYNSVALSRGTTTSLS--ATVESEG
BP19        302 IYIKAGVYREQVITPRKVNVMFMFGDGCATQITITPDRSWGLSPGTTTSLSGTVQVESEG
PER         261 VIYVKEGVYREITVWVPRDMAFVITLIGDPAKTRFTGSLMYADG--LLPMTATLGVNGEN
PEF1        279 VIYVKEGVYREITVWVAREMNVVITVIGDGPTRTKFTGSLMYADG--INTYKTAIFGVNGAN
PPE1        88  IIFIKAGVYKEYIDIPRSMTVWVLIGGPTKTRITGMSVVKDG--PTTFHTITVGVNGAN
AtPME6      281 IIRIKAGVYREKVTVTRKMPHVTFFIGDGNPNTLITGSLNFGIGK-VKPTPLTATITLEGDH
ATPME2      301 VHIKAGVYRENVEVTKKKNIMFLGDGRKTIITGSRNVVDG--STTFHSATVAVGER
ATPME3      311 VHIKAGVYRENVEVAKKKKNIMFMDGRTRTILITGSRNVVDG--STTFHSATVAVGER
PECS-1.1    303 IIRIKAGVYRENVEVTKKKNIMFMDGRTRTILITGSRNVVDG--STTFKSATVAVGER
PECS-1.2     80 IIRIKAGVYRENVEVTKKKNIMFMDGRTRTILITGSRNVVDG--STTFKSATVGGT---
PMEU1       302 VRIKAGVYRENVDVPPKKNIMFMDGKSNITLITSRNVVDG--STTFHSATVVRVCK
PECS-2.1    232 VIYIKAGTYRENLEVKLKN--HMFVGDGIGRTILITGSRNVVGG--ATTFKSATVAVGDN
LEPME1      156 VIYVKEGVYRENVEVSSNKNMLMIVGDGMVATITGSLNVVDG--STTFRSATLAAVGGG
LEPME2      267 VIYVKEGVYRENVEVSSKMKMLMIVGDGMHATITGSLNVVDG--STTFHSATLAAVKGG
LEPME3      262 VIYVKEGVYRENVEVTKKKNMLMIVGDGMNATITGSLNVVDG--STTFPSNLAAVGGG
RCPME1      271 VIYVKEGVYREKVEIKKKKNVMLVGDGMDATITGSLNMFIDG--TTTFHSATVAVGDG
AtPME7      194 VIYVKEGVYRENVEVTKKKNVMLVGDGIDVTVISGNSFIDG--WTTFRSATFAVSGRG
OsPME       333 VIYVKEGVYRENVKIGSKKTNMLVGDGAGTVVVGYSVHDM--YTTFHTATLAVAGAG
ATPME1      310 VIYVKEGVYRENVEVDSKKNVMLVGDGKGRITLITGSRNVVDG--PTTYETATFAVCKGG
PMEB        170 RARVNPHGQY@PADPAWYMYNACATKAGATINTTCS@VWWSQS---NDFQLKNITVWVAL

AtPME9      212 YIEGTVDYDFIFGRGASLYLMTQLHAVGDGLRVIAAHNR@STTE----QNGYSFVHCKRVTG-
AtPME10     212 YIEGTVDYDFIFGRGASLYLMTQLHAVGDGLRVITAQGR@SATE----QNGYTFVHCKRVTG-
AtPME8      193 YVEGTVDYDFIFGSGTSMYLGTOHVHVVDGIRVIAAHACKSAEE---KSGYSFVHCKRVTG-
ATPME4      419 VVSGTVDFIFGKSAITVIONTLIVVRKSGKQYNTVITADGNELGLGNKIGIVLNCRIVFD
BP19        417 VVSGTVDFIFGKSAITVIONSLILCRGSPGCTVHVITADGNEKGAVKIGIVLHNCRIIMAD
PER         374 SISGTTDMI@VGDFAVVFQNGKLVVRKPLEEQCFVADGRITK-S@SSSGYVFCSC@FTTGE
PEF1        392 AISGTTDFVFGDAFQV@ONKLVCFV@AKGQKCLVITAGGRDK-QNSABALVFLSS@FTTGE
PPE1        201 TITGTVDFIFGNGEAVL@ONCKVIVRKP@QNC@SCHVTAQGRTE-F@CKG@IVL@NCH@KPK
AtPME6      395 TVSGTVDFIFGDAKCL@ONCKVIVRKNPKGCT@CHVTAQGRSM-VRESTGLVLIHGCHITGD
ATPME2      414 HITGTVDFIFGNAAAVL@QCDIHARRPNSG@KNNVTAQGR@D-PNCNTGIVIQNCRIG@T
ATPME3      424 LIAGTVDFIFGNAAAVL@QCDIHARRPNSG@KNNVTAQGR@D-PNCNTGIVIQKCRIG@T
PECS-1.1    416 LIAGTVDFIFGNAAAVL@QCDIHARRPNSG@KNNVTAQGR@D-PNCNTGIVIQKCRIG@T
PECS-1.2    135 -----AAV@L@N@C@D@I@H@A@R@P@N@S@G@K@N@N@V@T@A@Q@G@R@D-PNCNTGIVIQKCRIG@T
PMEU1       415 LVAGTVDFIFGNGEAVV@QCDIHARRPNSG@KNNVTAQGR@D-PNCNTGIVIQKCRIG@T
PECS-2.1    343 DIYGTVDYDFIFGNAAAVL@QNCNIFARKE-PNRRITLIT@A@Q@G@R@D-PNCSTGLI@HNCR@V@T@A
LEPME1      269 YVTGTVDFIFGNAAAVV@QKCKLVARKPKGKY@QNNVTAQGR@D-PNCATGTSIQFCN@I@S
LEPME2      380 YVTGTVDFIFGNAAAVV@QKCKLVARKPKGKY@QNNVTAQGR@D-PNCATGTSIQFCN@I@S
LEPME3      374 YVTGTVDFIFGNAAAVV@QKCKLVARKPKR@KNNVTAQGR@D-PNCATGTSIQFCN@I@S
RCPME1      384 FITGTVDFIFGNAAAVV@QKSKLVARKPMSN@KNNVTAQGR@D-PNCNT@T@S@I@Q@C@N@V@I@P@S
AtPME7      307 TITGTVDFIFGDGTVV@QNCQILAKGLPNCKNTIT@A@Q@G@K@D-VNCP@S@I@Q@F@S@N@I@S@D
OsPME       446 DVAGTVDFIFGNAAAVL@QNCNIMARR@LP@G@C@N@V@T@A@Q@G@R@D-PNCSTG@S@V@H@G@C@R@L@P@S
ATPME1      423 DVTGTVDFIFGSAAVV@QGCKL@P@E@Q@L@S@N@C@N@T@I@A@Q@G@K@D-PNC@S@G@S@I@Q@C@I@S@N
PMEB        286 YIEGDVDYVFG@R@A@T@V@D@R@V@R@F@T@V@S@S@R@G@S@K@E@A@V@V@F@P@D@S@I@P--SVKY@G@L@V@I@N@S@C@I@T@G@D


AtPME9      267 -----VGTGIVLGRANM@SHP@VWVYSY@EMSSV@MPS@G@Q@C@N@R@V@R---A@H@K@I@V@F
AtPME10     267 -----TGTGIVLGRANM@SHP@VWVYAF@EMTSV@MPS@G@Q@C@N@L@N@R---G@Y@D@K@I@V@F
AtPME8      248 -----TGGGIVLGRANM@SHP@VWVYAY@EMTSV@MPS@G@Q@C@N@K@P---A@H@K@I@V@F
ATPME4      479 RKL@I@P@E-----RLTV@TYLGRPW@K@K@F@S@T@V@I@M@S@D@E@M@G@D@I@R@E@G@K@I@M@D@G@E---S@F@H@K@S@C@R
BP19        477 KELEAD----RLTVK@S@Y@LGRPW@K@P@F@T@I@A@V@I@G@T@E@I@G@D@L@I@C@P@T@G@M@E@M@Q@G@E---K@F@H@L@I@A@T
PER         433 PEWAKI----DP-KI@Y@V@LGRPW@K@S@YS@V@V@I@M@D@S@I@D@D@I@F@D@E@G@Y@M@P@M@M@G@S---A@F@K@D@I@C@T
PEF1        451 PALTSV----TP-KL@S@Y@LGRPW@K@L@YS@V@V@I@M@D@S@I@D@D@I@F@D@E@G@Y@M@P@M@V@G@G---A@F@K@D@I@C@T
PPE1        260 TDYFSL----SPPSRTY@LGRPW@K@E@YS@R@T@I@M@Q@S@I@D@K@F@E@P@E@G@A@P@M@I@T@N--F@G@R@D@S@Y
AtPME6      454 PAYIPM----KSVNKA@Y@LGRPW@K@E@F@S@R@T@I@M@K@T@I@D@D@V@I@D@P@A@G@L@P@M@S@G@D---F@A@L@K@T@L@Y
ATPME2      473 SDLLAV----KGTFFTY@LGRPW@K@E@YS@R@T@V@I@M@Q@S@I@D@S@V@I@R@E@G@M@E@M@S@G---S@F@A@L@D@T@L@T
ATPME3      483 SDLQSV----KCSFFTY@LGRPW@K@E@YS@R@T@V@I@M@Q@S@I@D@S@V@I@R@E@G@M@E@M@S@E@W@T@G---T@F@A@L@N@T@L@T
PECS-1.1    475 SDLKPV----QGSFFTY@LGRPW@K@E@YS@R@T@V@I@M@Q@S@I@D@L@I@H@P@A@G@M@E@M@D@G---N@F@A@L@N@T@L@F
PECS-1.2    181 SDLKPV----QGSFFTY@LGRPW@K@E@YS@R@T@V@I@M@Q@S@I@D@V@I@H@P@A@G@M@E@M@D@G---N@F@A@L@N@T@L@F
PMEU1       474 SDLRPW----QKSFFTY@LGRPW@K@E@YS@R@T@V@I@M@Q@S@I@D@V@I@Q@P@A@G@M@E@M@N@G---N@F@A@L@D@T@L@F
PECS-2.1    401 SDLKPV----QSSV@T@L@L@G@R@P@W@K@O@Y@S@R@T@V@I@K@I@F@I@D@S@L@I@H@P@A@G@M@E@M@S@G---D@F@A@L@N@T@L@Y
LEPME1      328 SDLEPW----LKEFFTY@LGRPW@K@E@YS@R@T@V@V@M@E@S@Y@L@G@L@I@H@P@A@G@M@E@M@D@G---D@F@A@L@K@T@L@Y
LEPME2      439 SDLEPW----LKEFFTY@LGRPW@K@K@YS@R@T@V@V@M@E@S@Y@L@G@L@I@H@P@A@G@M@E@M@D@G---D@F@A@L@K@T@L@Y
LEPME3      433 PDLEPW----MNEYRTY@LGRPW@K@K@S@R@T@V@V@M@Q@S@I@D@G@H@I@D@S@G@M@E@M@R@G---D@F@A@L@K@T@L@Y
RCPME1      443 SDLKPV----QGSIRTY@LGRPW@K@K@YS@R@T@V@V@L@Q@S@V@D@S@H@I@D@P@A@G@M@E@M@D@R@A-SKDFL@C@T@L@Y
AtPME7      366 ADLVPY----LNTT@TY@LGRPW@K@L@YS@R@T@V@I@R@N@M@S@D@V@I@R@E@G@L@E@M@N@A---D@F@A@L@D@T@L@F
OsPME       505 PELELAPAARRGRA@TY@LGRPW@K@P@YS@R@V@V@M@S@Y@D@A@G@V@H@A@G@L@A@D@A@S@G--R@P@D@I@L@Y
ATPME1      482 GNWIA@P-----TY@LGRPW@K@E@F@S@T@V@I@M@E@V@I@G@A@V@I@R@E@S@G@M@S@V@S@G---V@D@P@P@A@S@I@V
PMEB        344 N@Y@R@G@A-----Q@K@A@R@L@G@R@A@D@Q@G@A@Q@T@G@Y@L@P@G@K@T@A@N@G@Q@L@V@I@R@D@S@T@I@D@S---S

```

Isolation and study of a ubiquitously expressed tomato pectin methylesterase regulatory region

AtPME9	313	YGEYMC	TGPG	SHKAKR	VAH---	TQDIDN	KEASQ	FLTLG	VTKG	----	SKULL	PPPP	AY---
AtPME10	313	YGEYKCF	GPGE	SHLEKRV	PY---	TQDIDN	KEVTP	FLTLG	VTKG	----	STULL	PPPKY	---
AtPME8	294	YGEYKCS	GPGE	SHKAKR	VPPF---	TQDIDD	KEANCF	LSSLG	VIQG	----	SKULL	PPPP	AL---
ATPME4	532	YVEYNNR	RGPC	AFANRR	RVMMAK	VARSA	AAEVN	GF	TAANWL	GPIN	-----	WIQE	ANVP
BP19	530	YVEYNNR	RGPC	AMTAA	RRVPM	AKMAKS	AAEVE	RF	TVANWL	TPAN	-----	WIQE	ANVP
PER	485	FVEYNNK	GGP	GADT	NLRV	KWHG	VKVL	SNVA	AEY	TPGK	FE	IVN	ATARD
PEF1	503	FVEYNNK	GGP	GADT	NLRV	KWHG	VKVL	SNVA	AEY	TPGK	FE	IVN	ATARD
PPE1	314	YREYQNR	GP	GAAL	DRIT	TKGF	QKGF	TGE	AAQK	FTAG	VY	INN	----
AtPME6	507	YAEHMM	TGPG	ENQA	CRV	KMPG	IK	KLTP	-QD	ALL	VTG	DR	LRG
ATPME2	526	YREYLMR	GGG	AGTAN	RVKK	GYR	VIT	SD	TE	IQPF	TAG	CF	IGG
ATPME3	536	YREYSMT	GAG	AGTAN	RVKK	GYR	VIT	AAEA	QK	FTAG	CF	IGG	----
PECS-1.1	528	YGEHQN	SGG	AGT	SRV	KMG	FR	VIT	S	ATE	AQ	FT	PGSF
PECS-1.2	234	YGEHQN	SGG	AGT	SRV	KMG	FR	VIT	S	ATE	AQ	FT	PGSF
PMEU1	527	YGEYANT	GAG	APTS	GRV	KMG	FR	VIT	S	TE	AQ	FT	PGSF
PECS-2.1	454	YREYMMT	GPGE	STAN	RVKK	GYH	VLT	SPS	QVS	QF	TV	EM	TAG
LEPME1	381	YGEFMMN	GGG	AGT	SRV	KMPG	YH	VIT	DP	AK	MP	FT	VAKL
LEPME2	492	YGEFMMN	GGG	AGT	SRV	KMPG	YH	VIT	DP	AK	MP	FT	VAKL
LEPME3	486	YGEFMMN	GGG	AGT	SRV	KMPG	YH	VIT	DP	AK	MP	FT	VAKL
RCPME1	498	YGEYLSN	GGG	AGT	SRV	TK	YH	I	K	T	A	E	AS
AtPME7	419	YGEYNNY	GPGE	GLSS	RVKK	GYH	VFN	NSD	Q	ANN	F	TV	SQ
OsPME	563	YGEYRNS	GGG	AAV	GRV	PPG	HR	V	TKL	PE	E	ME	F
ATPME1	531	YGEYKNT	GPGE	DVTC	RVKK	AGY	KP	MSD	AE	AK	F	TV	TL
PMEB	388	YDLANP	WGA	LLMT	DR	PF	EG	-----	NIS	PQR	DL	DI	HN
AtPME9													
AtPME10													
AtPME8													
ATPME4	585	GL											
BP19	583	GL											
PER	545	ALD	ATS	NQ	GAT	PG	Q	GT	V	T	G	G	A
PEF1	563	CPM											
PPE1	370	GMM	KV										
AtPME6	561	KV											
ATPME2	581	SL											
ATPME3	591	GL											
PECS-1.1	583	GL											
PECS-1.2	289	GL											
PMEU1	582	GL											
PECS-2.1	509	GL											
LEPME1	436	GLY	D										
LEPME2	547	GLY	D										
LEPME3	541	GLV	E										
RCPME1	553	GL											
AtPME7	474	GLY	I										
OsPME	618	GLT	V										
ATPME1	586	S											
PMEB	434												

**Figure 3. Aminoacid Alignment of 22 Plant PME's and *Erwinia chrysanthemi* PME.** PMEU1 (U49330), LEPME1 (U70677), LEPME2 (U70675) and LEPME3 (U70676) are from *Lycopersicon esculentum*; AtPME1 (NP\_175787), AtPME2 (PC4168), AtPME3 (NP\_188048), AtPME4 (AF077855), AtPME6 (AAF63815), AtPME7 (T05202), AtPME8 (NP\_568181), AtPME9 (NP\_196359) and AtPME10 (NP\_196360) are from *Arabidopsis thaliana*; PER (AJ249611) and PEF1 (AJ249611) are from *Medicago truncatula*; PECS-1.1 (U82973), PECS-1.2 (U82974) and PECS-2.1 (U82975) are from *Citrus sinensis*; Bp19 (X56195) is from *Brassica napus*, PpE1 (L27101) is from *Petunia inflata*, RCPME1 (AF081457) is from *Pisum sativum*, OsPME1 (BAA96597) is from *Oriza sativa* and PME B (X84665) is from *Erwinia chysanthemi*. Alignment of deduced aminoacid was done using GCG's Pileup Program (Genetics Computer Group, Madison, WI).



**Figure 4. Phylogenetic Analysis of 22 Plant PME's and *Erwinia chrysanthemii* PME.** PMEU1 (U49330), LePME1 (U70677), LePME2 (U70675) and LePME3 (U70676) are from *Lycopersicon esculentum*; AtPME1 (NP\_175787), AtPME2 (PC4168), AtPME3 (NP\_188048), AtPME4 (AF077855), AtPME6 (AAF63815), AtPME7 (T05202), AtPME8 (NP\_568181), AtPME9 (NP\_196359) and AtPME10 (NP\_196360) are from *Arabidopsis thaliana*; PER (AJ249611) and PEF1 (AJ249611) are from *Medicago truncatula*; PECS-1.1 (U82973), PECS-1.2 (U82974) and PECS-2.1 (U82975) are from *Citrus sinensis*; Bp19 (X56195) is from *Brassica napus*, PpE1 (L27101) is from *Petunia inflata*, RCPME1 (AF081457) is from *Pisum sativum*, OsPME1 (BAA96597) is from *Oriza sativa* and PME1 (X84665) is from *Erwinia chrysanthemii*. Numbers are the bootstrap values. Phylogenetic analysis were done using PHYLIP (phylogeny inference package) ver 3.5c.




```

-847 TTAGATATTTAGTTTGTATCATAGATTCTCAATTTTTGAAGAAAAGTAGCCGCT
-794 CGTCTCCGGTTCTTGAAGAACTTTACTTATAGCTAATTTAGAGAGCTTTACGT
-741 AAGATATGAGGTTCTGCTCCTCCAAGGTTCCAAAGTCCACCCTTACATATATC
-688 GATTTTATAGATAGATTAAATGGGAAATAATACCACAAAAGAAGTAAACAATTG
-635 GATTCTTGTAATTTAGCCTACACACTAAACTACTCCATAATTAGCAGTTCAT
-582 TAAAAACACTTAATTAGGCCAAAAACAAGAAAGCTAGTTAACGAACATCTGCCAGG
-529 AAGTGTGATGCTAAAAAACAATTAATTACTAGGAAAAATAGACTAGGCATTA
-476 ACATTAACATCTAGTTAAATTTATCTATTTTACGATGTCCAATATCCTGGCCG
-423 CCCAGGCTAAAAGTTTTACTTCTTTTTGTTGCTTCACCCACTTGTTATTCTGTG
-370 AACAAAATTC AATATACAGCTAATCATGTCACTTTCAAATATCAATGATCGG
-317 AGTGATAATAATTGATTAAGTTTTAGATAAAGGCACGACTCAGCACAACAAGA
-264 ATTCTTTTCAAAAAGTAATCAAACAAGAAAAGTATCGATACCTTCCCACCAT
-211 TCACGTGGCATCTATTATTACATGCAAAATTCACATTATGACCAAAAACAATAAA
      IV
-158 TCAAAATTTATTCCTAGTAAATGCCATACTGTACTTTTTAAAAAATTCGAATAT
-105 TATATTTCAAAAAATTAGCGTATATATAGTTTTTCGACAACTTGATAATATAT
-52 ATATAAAAACATAAAATTTAAAAGTAAAAACA AAAATAGAACGTTGAAATTA

```


**Figure 5. DNA sequence of the PMEU1 Promoter.** Shown are the longest direct repeats (numbered in bold), mirror repeats (arrows in opposite directions), putative TATA box (doubled underlined), translation start site (bold) and putative cis-acting elements (roman numbered and boxed). The software used to find the promoter characteristics is explained in the body of the paper.



**Figure 6. Chimeric PMEU1 promoter constructs and average of GUS activity in the transgenic tobacco plants.**

A. The chimeric constructs used in plant transformation. Numbers below the shadowed bar are indicating the size of the *pmeu1* promoter in each construct. Arrow is indicating the translation start site for the *pmeu1* transcribed region.

B. Average of GUS activity from leaf of about 50 tobacco transgenic plants analyzed. Shown are the average and standard deviation values. Differences in GUS activity levels among all three constructs were statistically significant ( $p < 0.05$ ).



**Figure 7. Average of GUS activity for root, stem and leaf of tobacco transgenic plants.** Six independent transgenic tobacco plants harboring each of the three constructs were used to determine the average of GUS activity in root, stem and leaf. Shaded, white and black bars are the average of GUS activity for plants harboring 0.267 kb of promoter size, 1.306 kb of promoter and 2.59 kb, respectively. Lines in bars are indicating the standard deviation. Both average and standard deviation values were calculated by transforming back the square root transformed data used in the statistical analysis. Root and leaf values are statistically significant ( $p < 0.05$ ). GUS activity for the constructs including 1.306 kb and 2.59 kb of promoter size showed significant differences when comparing root with leaf and stem. For the construct including 0.267 kb of promoter size, statistical analysis did not detect differences ( $p > 0.05$ ).